

RESEARCH ARTICLE

Open Access



Genetic analysis of enteropathogenic *Escherichia coli* (EPEC) adherence factor (EAF) plasmid reveals a new deletion within the EAF probe sequence among O119 typical EPEC strains

Nathalia B. Teixeira¹, Thais C. G. Rojas², Wanderley D. da Silveira², Cecília Matheus-Guimarães¹, Neusa P. Silva³ and Isabel C. A. Scaletsky^{1*}

Abstract

Background: Enteropathogenic *Escherichia coli* (EPEC) are classified into typical and atypical strains based on the presence of the *E. coli* adherence factor (EAF) plasmid. The EAF plasmid contains the *bfp* (bundle-forming pilus) operon and the *perABC* (plasmid encoded regulator) gene cluster. A 1-kb cryptic region of EAF plasmid has been widely used as a genetic probe for EPEC detection. However, some EPEC strains may harbor an EAF plasmid lacking the EAF probe sequence, which makes the differentiation between typical and atypical a complex task. In this study, we report the genetic analysis of the EAF plasmid-encoded genes in a collection of EPEC clinical isolates.

Methods: A total of 222 EPEC clinical isolates, which were previously classified as typical ($n = 70$) or atypical ($n = 152$) by EAF probe reactivity, were screened for the presence of different EAF sequences by PCR and DNA hybridization.

Results: All typical strains possessed intact *bfpA* and *perA* genes, and most of them were positive in the PCR for EAF probe sequence. However, a subset of 30 typical strains, 22 of which belonged to O119 serogroup, presented a 1652 pb deletion in the region between 1093-bp downstream *perC* and 616-bp of the EAF fragment. The *bfpA*, *bfpG*, and *per* genes were found in all typical strains. In addition, 32 (21 %) atypical strains presented the *perA* gene, and 20 (13.2 %) also presented the *bfpA* gene. Among the 32 strains, 16 belonged to the O119:H2, O119:HND, and ONT:HND serotypes. All 32 atypical strains contained *perA* mutation frameshifts and possessed an *IS1294* element upstream of the *per* operon as detected by PCR followed by restriction fragment length polymorphism (RFLP) typing and multiplex PCR. Among the 20 *bfpA* probe-positive strains, eight O119 strains possessed deletion in the *bfp* operon at the 3'end of *bfpA* due to an *IS66* element.

Conclusion: Our data show that typical O119 strains may contain a deletion within the EAF probe sequence not previously reported. This new finding suggests that care should be taken when using the previously described EAF PCR assay in epidemiological studies for the detection of typical O119 strains. In addition, we were able to confirm that some atypical strains carry vestiges of the EAF plasmid.

Keywords: Enteropathogenic *Escherichia coli*, EAF plasmid

* Correspondence: scaletskyunifesp@gmail.com

¹Departamento de Microbiologia, Imunologia e Parasitologia, Universidade Federal de São Paulo, Rua Botucatu, 862, 3 andar, São Paulo 04023-062 São Paulo, Brazil

Full list of author information is available at the end of the article

Background

Enteropathogenic *Escherichia coli* (EPEC) are an important cause of infantile diarrhea in developing countries, particularly in Brazil [1–3]. EPEC strains produce a characteristic intestinal histopathology called the attaching-and-effacing (A/E) lesion, which is characterized by microvilli effacement and intimate bacterial adherence to the epithelial membrane. The genes responsible for A/E lesion phenotype are encoded on 35.6-kb chromosomal pathogenicity island known as the locus of enterocyte effacement (LEE). The LEE island comprises approximately 40 genes and encodes the components of a type III secretion system, various effector molecules, and the intimin adhesin which is encoded by the *eae* gene [4, 5].

EPEC strains are classified into typical and atypical based on the presence of the large virulence EPEC adherence factor (EAF) plasmid [6]. The EAF plasmid encodes a type IV fimbria known as the bundle-forming pilus (BFP) that mediate localized adherence (LA) to epithelial cells [7–9]. A 14-gene operon is necessary for BFP production [10], with *bfpA* encoding the major structural subunit (bundlin). A second operon on the EAF plasmid is the plasmid-encoded regulator (Per), consisting of three genes (*perA*, *perB*, and *perC* genes) [11], which activates genes within the LEE and the *bfp* operon [12–14].

Additional criteria for classifying isolates as EPEC include the detection of specific serogroups [6]. Classic EPEC O serogroups include O55, O86, O111, O114, O119, O125, O126, O127, O128, and O142. By multilocus enzyme electrophoresis analysis of allelic differences among housekeeping genes, typical EPEC strains have been subtyped into two major lineages, previously designated EPEC1 and EPEC2 [15, 16]. The EPEC1 includes widespread serotypes such as O55:H6 and O119:H6, whereas EPEC2 consists of serotypes with more limited occurrence such as O111:H2 and O114:H2. Based on a whole-genome phylogeny and analysis of type III secretion system effectors, typical EPEC strains have been demonstrated to cluster in three main lineages, designated EPEC1, EPEC2, and EPEC4 [17]. According to the phylogenomic analyses by Hazen et al. [17] the term atypical EPEC refers to a group of phylogenetically diverse isolates than often are more similar to *E. coli* of other pathovars than EPEC. However, although the term atypical EPEC may be considered misleading, atypical strains are identified by the presence of *eae* and the absence of the EAF probe sequence as well as the Shiga toxin-encoded genes [6].

A 1-kb cryptic region of EAF plasmid has been widely used as a genetic probe for EPEC detection [18]. However, some EPEC strains, as demonstrated by BFP production, may in fact harbor an EAF plasmid lacking the EAF probe sequence [19, 20], which makes the differentiation

between typical and atypical complex [21]. Other EPEC strains harbor an EAF plasmid that shares a conserved backbone, and is in many ways similar to pMAR7 plasmid of EPEC1 strain E2348/69 and pB171 plasmid of EPEC2 strain B171, but with inactivating deletions in the *bfp* and *per* operons that are required for LA [22, 23]. Such strains are phenotypically “atypical”, since they do not produce a typical LA pattern on epithelial cells, even though they carry probe-detectable EAF plasmids. In this study, we report the genetic analysis of the EAF plasmid-encoded genes in a collection of EPEC clinical isolates.

Results and discussion

In this study, we analyzed a collection of 222 EPEC clinical isolates, including strains of the classic and nonclassic EPEC serotypes, which were previously classified as typical ($n = 70$) or atypical ($n = 152$) by EAF probe reactivity. Initially, all 222 strains were screened for the EAF probe sequence by using a PCR with primers located 100 nucleotides upstream (5'-CGCCATTTATTTTAAGACGAACA) and 82 nucleotides downstream (5'-CGCTTCTGCTTTTGACGG) the EAF sequence of pMAR2 [24]. As shown in Tables 1 and 2, the majority of typical strains yielded the expected 1260-bp amplicon. Interestingly no PCR product was obtained from a subset of 30 typical strains, 22 of which belonged to O119 serogroup. To verify this, a new primer located 20 nucleotides upstream *perB* (5'-GAGCACTCGAAATGAA GAAC) was designed to include the region between *perBC* and EAF sequence. All 30 typical strains showed a PCR product approximately 1.7-kb smaller than the expected (3.9 kb). DNA sequencing revealed the presence of a 1652 pb deletion not previously reported, in the region between 1093-bp downstream *perC* and 616-bp of the EAF fragment. As expected all the atypical strains were PCR-negative.

As shown in Tables 1 and 2, the *bfpA*, *bfpG*, and *per* genes, detected by PCR and colony blot hybridization, were present in all typical strains. In addition, 32 (21 %) atypical strains presented the *per* genes, and 20 (13.2 %) also presented the *bfpA* gene. Among the 32 strains, 18 strains belonged to the serotypes O119:H2, O119:HND, O142:H2, and ONT:HND atypical strains. Although 20 of 32 strains hybridized with the *bfpA* probe, *bfpA* could not be amplified by PCR from eight O119 strains, suggesting that the 3' end of the gene was deleted and replaced with an IS66-like element as described by Bortolini et al. [22]. To verify this, we carried out PCR with primers targeting the 5' end of *bfpA* and the IS66-like element previously described [22]. All the eight O119 strains yielded the expected amplicon, and DNA sequencing confirmed the presence of a 1053-bp IS element inserted into *bfpA* at position 262 with significant similarity to IS66 (97 %). We evaluated the level of

Table 1 Results of PCR and hybridization for EAF-encoded genes among typical EPEC strains

Clonal group	Serotype	No. of strains	Source	Presence or absence of:				IS element upstream of <i>per</i>	Presence or absence of:					<i>perBC</i> -EAF PCR product
				<i>bfpA</i> (probe)	<i>bfpA</i> (PCR)	<i>bfpG</i> (PCR)	<i>trcP</i> (probe)		<i>per</i> (probe)	<i>perA</i> (PCR)	EAF (PCR)	<i>orf35-36</i> (probe)	<i>orf61-62</i> (probe)	
EPEC1	O55:H6, NM	13	Patients	+	+	+	–	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC1	O55:NM	1	Control	+	+	+	–	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC1	O86:NM, H34	2	Patients	+	+	+	+	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC2	O111:NM	5	Patients	+	+	+	+	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC2	O111:H2	4	Patients	+	+	+	+	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC1	O119:H6, NM	16	Patients	+	+	+	–	IS1(vξ)	+	+	–	+	–	2.2-kb
EPEC1	O119:H6	1	Patient	–	+	+	–	IS1(vξ)	–	+	–	+	–	2.2-kb
EPEC1	O119:H6, NM	5	Patients	+	+	+	–	IS1(vξ)	+	+	–	–	+	2.2-kb
EPEC1	O119:H6	2	Controls	+	+	+	+	IS1(vξ)	+	+	+	+	–	3.9-kb
EPEC1	O119:H6	1	Control	+	+	+	+	IS1(vξ)	–	+	+	+	–	3.9-kb
EPEC1	O127:H6, NM	3	Controls	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
EPEC1	O127:NM	1	Patient	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
Unknown	O2:H2, H45	2	Patients	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
Unknown	O101:H33	1	Control	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
Unknown	O145:HNT	1	Control	+	+	+	–	IS1(vξ)	+	+	–	+	–	2.2-kb
Unknown	O162:NM, H33	2	Patients	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
Unknown	O157:HND	2	Patients	+	+	+	–	IS1(vξ)	+	+	–	+	–	2.2-kb
Unknown	ONT:H45; HND	3	Patients	+	+	+	–	IS1(vξ)	+	+	+	+	–	3.9-kb
Unknown	ONT:HND	5	Patients	+	+	+	–	IS1(vξ)	+	+	–	+	–	2.2-kb
EPEC1	E2348/69 (O127:H6)			+	+	+	–	IS1(vξ)	+	+	+	+	+	3.9-kb
EPEC2	B171 (O111:NM)			+	+	+	+	IS1(vξ)	+	+	+	+	+	3.9-kb

NM Nonmotile; ND Nondetermined; NT Nontypeable

Table 2 Characteristics of aEPEC strains positive for EAF plasmid-encoded genes either by PCR or hybridization assay

Strain	Serotype	Source	Presence or absence of:				IS element upstream of <i>per</i>	Presence or absence of:					HEp-2 adhesion pattern
			<i>bfpA</i> (probe)	<i>bfpA</i> (PCR)	<i>bfpA-IS66</i>	<i>trcP</i> (probe)		<i>per</i> (probe)	<i>perA</i> (PCR)	EAF (PCR)	<i>orf35-36</i> (probe)	<i>orf61-62</i> (probe)	
A103	O9:HND	Patient	-	-	-	+	<i>IS1294</i>	+	+	-	-	-	-
A148	O37:NM	Patient	-	-	-	-	<i>IS1294</i>	+	+	-	-	-	-
A141	O49:HND	Patient	-	-	-	-	<i>IS1294</i>	+	+	-	+	-	-
A69	O55:NM	Patient	+	+	-	+	<i>IS1294</i>	+	+	-	-	-	LAL
A152	O96:NM	Control	-	-	-	-	<i>IS1294</i>	+	+	-	-	-	-
A144	O98:HND	Patient	-	-	-	+	<i>IS1294</i>	+	+	-	+	-	-
A140	O108:HND	Control	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A60	O119:H2	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A62	O119:H2	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A66	O119:H2	Control	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A67	O119:H2	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A75	O119:HND	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A90	O119:HND	Patient	+	+	-	-	<i>IS1294</i>	+	+	-	-	-	LA
A111	O119:HND	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A127	O119:HND	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A131	O119:HND	Patient	+	-	+	-	<i>IS1294</i>	+	+	-	-	-	LAL
A97	O128:HND	Patient	-	-	-	-	<i>IS1294</i>	+	+	-	-	-	-
A126	O128:NM	Patient	-	-	-	-	<i>IS1294</i>	+	+	-	-	-	-
A11	O142:NM	Patient	+	+	-	+	<i>IS1294</i>	+	+	-	+	-	LA
A129	O142:HND	Patient	+	+	-	+	<i>IS1294</i>	+	+	-	+	-	LAL
A124	O157:NM	Patient	-	-	-	-	<i>IS1294</i>	+	+	-	-	-	-
A95	ONT:H21	Patient	+	+	-	+	<i>IS1294</i>	+	+	-	-	-	LAL
A65	ONT:NM	Control	+	+	-	-	<i>IS1294</i>	+	+	-	-	-	LAL
A74	ONT:NM	Patient	+	+	-	-	<i>IS1294</i>	+	+	-	-	-	LAL
A63	ONT:HND	Patient	+	+	-	-	<i>IS1294</i>	+	+	-	+	+	LAL
A64	ONT:HND	Patient	+	+	-	-	<i>IS1294</i>	+	+	-	+	+	LAL
A78	ONT:HND	Patient	+	+	-	-	<i>IS1294</i>	+	+	-	+	-	LAL
A146	ONT:HND	Control	+	+	-	-	<i>IS1294</i>	+	+	-	+	-	LAL

Table 2 Characteristics of aEPEC strains positive for EAF plasmid-encoded genes either by PCR or hybridization assay (Continued)

A57	ONT:HND	Patient	–	–	–	–	<i>IS1294</i>	+	+	–	–	–	–
A16	ONT:HND	Control	–	–	–	–	<i>IS1294</i>	+	+	–	+	–	–
A118	ONT:HND	Patient	–	–	–	–	<i>IS1294</i>	+	+	–	–	–	–
A136	ONT:HND	Patient	–	–	–	–	<i>IS1294</i>	+	+	–	+	–	LAL

ND Nondetermined; NM Nonmotile; LA Localized adherence patter; LAL LA-like pattern

expression of *bfpA* by RT-PCR in the 20 *bfpA*-positive strains. The RT-PCR results showed that the *bfpA* gene was transcribed only by the strains carrying the intact *bfpA* gene sequence (Fig. 1). Two of these strains, A90 (O119:HND) and A11 (O142:NM), present the LA pattern. Considering the BFP production as a truly phenotype marker of typical EPEC strains, these strains are in fact typical EPEC rather than atypical EPEC as originally classified by the EAF-probe reactivity.

The EAF plasmids from two well-studied EPEC strains have been sequenced [25, 26]. The major difference between pMAR7 (E2348/69, EPEC1 lineage) and pB171 (B171, EPEC2 lineage) is the presence of conjugative transfer (*tra*) genes on pMAR7, absent in pB171 [26]. Apart from the *tra* region, several other ORFs are present in pB171 but not in pMAR7, such as the region *orf35–36* (truncated homolog of the EHEC *tox*B gene), the region *orf61–62* (*gadB* [truncated] and *gadC* homologs), and the putative chaperonine *trcP* located between the *bfp* and *per* operons flanked by insertion (IS) elements. As shown in Tables 1 and 2, a subset of typical strains O55, O86, O111 and O119 strains were positive for one or more pB171 derived-probes, while most of atypical EPEC strains did not hybridize with one or more of the probes (Fig. 2).

The sequence of plasmid pB171 reveals that a partial *IS1*(v ξ) homolog lies immediately upstream of the *per* operon [26]. This *IS1*(v ξ) element is replaced by a partial *IS1294* in O119:H2 (MB80), O128:H2, and O142:H6 (O142#15) strains containing *perA* mutation frameshifts, leading to premature truncation and consequent inactivation of the gene [23]. Okeke et al. [23] developed a multiplex PCR which gives a 650-bp product in strains with the *IS1*(v ξ) element and a 500-bp product in strains that have the *IS1294*, and a *SspI*-based PCR-RFLP typing which distinguishes normal *perA* alleles from those with O119:H2, O128:H2, or O142:H6 type-frameshift. As

shown in Tables 1 and 2, all typical strains yielded a 650-bp product, while all 32 atypical strains with *perA* mutation frameshifts yielded a 500-bp product. In addition, all typical strains showed a *perA* RFLP pattern similar to that of control EPEC strain (E2348/69), while all 32 *perA* probe-positive atypical strains produced a pattern consistent with that of the MB80. The entire *perA* gene was amplified and sequenced in all 32 atypical strains, and DNA sequencing confirmed the presence of *perA* mutation identical to that of MB80 strain. These results suggest that the presence of an inactive *perA* gene could be used to differentiate typical from atypical strains. Interestingly, most strains carrying vestiges of EAF plasmid belong to the same serogroups as typical EPEC, suggesting that these strains may constitute a group of EPEC strains that carry a defective EAF plasmid rather than atypical EPEC isolates.

Conclusion

Our data show that typical O119 strains, which were the most prevalent isolates in São Paulo, in the past [27], may contain a deletion within the EAF probe sequence not previously reported. This new finding suggests that care should be taken when using the previously described EAF PCR assay [28] in epidemiological studies for the detection of typical O119 strains. In addition, we were able to confirm that some atypical strains carry vestiges of the EAF plasmid [29–31].

Methods

Strains

The strains examined in this report were isolated during epidemiological studies of acute diarrhea in children <2 years of age conducted in different regions of Brazil in 1999–2009 [2, 32–34]. These strains were identified by colony hybridization with *eae* and/or EAF probe sequences and serotyped, and most of them had also been characterized by the presence of LEE-associated DNA sequences, and adherence to HEp-2 cells [34, 35]. Strains were grown overnight at 37 °C in 5 ml of Luria-Bertani broth with shaking. Genomic DNA was isolated using the HiYield Genomic DNA Mini kit (Real Biotech Corporation, Taiwan), and was used as the template for the PCR assays.

Ethics statement

The study was approved by the ethics committee of the Universidade Federal de São Paulo, Brazil. Stool samples were obtained with the written informed consent from the parents or guardians of the children.

Screening for EAF plasmid genes

The presence of EAF plasmid-encoded genes was determined by colony hybridization and PCR. Colony





Fig. 1 Agarose gel electrophoresis of the RT-PCR products of O119 and O142 atypical EPEC strains. Lane1, 1-kb ladder plus (Gibco, BRL); lane 2, E2348/69 strain (positive control); lane 3, A60 strain; lane 4, A62 strain; lane 5, A66 strain; lane 6, A67 strain; lane 7, A75 strain; lane 8, A90 strain; lane 9, A111 strain; lane 10, A127 strain; lane 11, A131; lane 12, A11 strain; and lane 13, A129 strain



hybridizations were performed under high-stringency conditions at 65 °C employing probes that were labeled by random priming with [α - 32 P] dCTP as described previously [32]. Fragments probes for *bfpA* and *perABC*, were prepared from plasmid clones as described previously [22]. Probes for *bfpG*, *trcP*, *orf35-36*, and *orf61-62*, were prepared by PCR primers as described [23] using strain B171 as the template. The DNA fragments were purified, labeled with [α - 32 P] dCTP with a DNA labeling kit (Amersham Pharmacia Biotech Inc., EUA) and used as probes. Blots were hybridized in a solution containing the labeled probe (10^5 cpm), $5 \times$ standard saline citrate (SSC), $2 \times$ Denhardt's solution (Invitrogen), 0.1 % sodium dodecyl sulfate (SDS), and 5 mg/ml of salmon sperm DNA for 16 h at 65 °C. After hybridization, washes were done in aqueous solution with $2 \times$ SSC with 0.1 % SDS and exposed to X-ray film.

The *bfpA* gene was amplified by PCR with primers *bfpA*_114F (GTCTGCGTCTGATTCCAATA) and *bfpA*_521R (TCAGCAGGAGTAATAGC) as previously described [36].

The entire *perA* gene was amplified by PCR with primers K1693 (CCCAAGCTTTGGCAATGTTTCCTTG TGT) [23] and *perA*-24F (AACAAACGCGCATGAA GGTG) [22]. The 770-bp amplicon from *perA*-positive strains was digested with the restriction enzyme *SspI*, and restriction fragment length polymorphism (RFLP) analysis was performed by agarose gel electrophoresis.

The *per* upstream region to *IS1(vξ)* was amplified by PCR with primers K1547 (TGAGTCACCTCTGCCT GAG) and K1549 (TGGATTCTATTGTGTATTCGG), and the *per* upstream region to *IS1294* was amplified by PCR with primers K1978 (TGTGAGAGCTTCTCAGCA)

and K1549 (TGGATTCTATTGTGTATTCGG) as previously described [23].

RNA extraction and RT-PCR assays

Total RNA was extracted after bacterial growth in LB broth for 18 h at 37 °C with the RNase Mini extraction kit (Qiagen) according to the manufacturer's instructions. After extraction, approximately 1 μ g of total RNA was digested with DNase I (Qiagen) for 30 min at 37 °C, and the enzyme was then inactivated by adding 1 μ l of 25 mM EDTA and heating the solution at 65 °C for 10 min. To obtain the cDNA, the *SperScript* III One Step RT-PCR System with Platinum *Taq* DNA polymerase (Invitrogen) was used according to the manufacturer's specifications. Primers for 16S ribosomal protein were used to control PCR [37], and the assay was then carried out with primers *bfpA*_114F (GTCTGCGTCT GATTCCAATA) and *bfpA*_521R (TCAGCAGGAGTA ATAGC) as previously described [36]. PCR products were analyzed by 1 % agarose gel electrophoresis.

DNA sequencing

Nucleotide sequencing of the PCR products was performed at the Centro de Estudos do Genoma Humano-USP, São Paulo. Nucleotide sequence data were analyzed using SeqMan and MegAlign software and the BLAST tool (<http://www.ncbi.nlm.nih.gov/BLAST>).

Nucleotide sequence and accession number

The DNA sequences for the EAF region of typical EPEC O119 strains (T60 and T40) are available in NCBI database under accession numbers KT595240 and KT819171.

Abbreviations

EPEC: Enteropathogenic *Escherichia coli*; EAF: *E. coli* adherence factor plasmid; Bfp: Bundle-forming pilus; Per: Plasmid-encoded regulator.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NBT and CMG performed experiments. TCGR and WDS analyzed data and contributed to the preparation of the manuscript. ICAS designed the experimental procedures, supervised all the experimental works and wrote the manuscript. NPS participated in the design of the experiments and reviewed the paper. All authors read and approved the final manuscript.

Authors' information

Not applicable.

Acknowledgments

This study was supported by research grants from Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

Author details

¹Departamento de Microbiologia, Imunologia e Parasitologia, Universidade Federal de São Paulo, Rua Botucatu, 862, 3 andar, São Paulo 04023-062 São Paulo, Brazil. ²Departamento de Genética, Evolução e Bioagentes, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, Brazil. ³Disciplina de Reumatologia, Universidade Federal de São Paulo, São Paulo, Brazil.

Received: 4 May 2015 Accepted: 29 September 2015

Published online: 05 October 2015

References

- Levine MM, Edelman R. Enteropathogenic *Escherichia coli* of classic serotypes associated with infant diarrhea: epidemiology and pathogenesis. *Epidemiol Rev.* 1984;6:31–51.
- Araújo JM, Tabarelli GF, Aranda KR, Fabbriotti SH, Fagundes-Neto U, Scaletsky ICA. Typical enteroaggregative and atypical enteropathogenic types of *Escherichia coli* are the most prevalent diarrhea-associated pathotypes among Brazilian children. *J Clin Microbiol.* 2007;45:3396–9.
- Bueris V, Sircili MP, Taddei CR, Santos MF, Franzolin MR, Martinez MB, et al. Detection of diarrheagenic *Escherichia coli* from children with and without diarrhea in Salvador, Bahia, Brazil. *Mem Inst Oswaldo Cruz.* 2007;102:839–44.
- Finlay BB, Rosenshine I, Donnenberg MS, Kaper JB. Cytoskeletal composition of attaching and effacing lesions associated with enteropathogenic *Escherichia coli* adherence to HeLa cells. *Infect Immun.* 1992;60:2541–3.
- McDaniel TK, Jarvis KG, Donnenberg MS, Kaper JB. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc Natl Acad Sci U S A.* 1995;92:1664–8.
- Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev.* 1998;11:142–201.
- Donnenberg MS, Girón JA, Nataro JP, Kaper JB. A plasmid-encoded type IV fimbrial gene of enteropathogenic *Escherichia coli* associated with localized adherence. *Mol Microbiol.* 1992;6:3427–37.
- Girón JA, Ho AS, Schoolnik GK. An inducible bundle-forming pilus of enteropathogenic *Escherichia coli*. *Science.* 1991;254:710–3.
- Knutton S, Shaw RK, Anantha RP, Donnenberg MS, Zoragani AA. The type IV bundle forming pilus of enteropathogenic *Escherichia coli* undergoes dramatic alterations in structure associated with bacterial adherence, aggregation and dispersal. *Mol Microbiol.* 1999;33:499–509.
- Stone KD, Zhang HZ, Carlson LK, Donnenberg MS. A cluster of fourteen genes from enteropathogenic *Escherichia coli* is sufficient for the biogenesis of a type IV pilus. *Mol Microbiol.* 1996;20:325–37.
- Gómez-Duarte OG, Kaper JB. A plasmid-encoded regulatory region activates chromosomal *eaeA* expression in enteropathogenic *Escherichia coli*. *Infect Immun.* 1995;63:1767–76.
- Elliott SJ, Sperandio V, Girón JA, Shin S, Mellies JL, Wainwright L, et al. The locus of enterocyte effacement (LEE)-encoded regulator controls expression of both LEE- and non-LEE-encoded virulence factors in enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect Immun.* 2000;68:6115–26.
- Mellies JL, Elliott SJ, Sperandio V, Donnenberg MS, Kaper JB. The Per regulon of enteropathogenic *Escherichia coli*: identification of a regulatory cascade and a novel transcriptional activator, the locus of enterocyte effacement (LEE)-encoded regulator (Ler). *Mol Microbiol.* 1999;33:296–306.
- Tobe T, Schoolnik GK, Soheli I, Bustamante VH, Puente JL. Cloning and characterization of *bfpTWW* genes required for the transcriptional activation of *bfpA* in enteropathogenic *Escherichia coli*. *Mol Microbiol.* 1996;21:963–75.
- Ørskov F, Whittam TS, Cravioto A, Ørskov I. Clonal relationships among classic enteropathogenic *Escherichia coli* (EPEC) belong to different O groups. *J Infect Dis.* 1990;162:76–81.
- Whittam TS, McGraw E. Clonal analysis of EPEC serogroups. *Rev Microbiol.* 1996;27:7–16.
- Hazen TH, Sahl JW, Fraser CM, Donnenberg MS, Scheut F, Rasko DA. Refining the pathovar paradigm via phylogenomics of the attaching and effacing *Escherichia coli*. *PNAS.* 2013;110:12810–5.
- Nataro JP, Baldini MM, Kaper JB, Black RE, Bravo N, Levine MM. Detection of an adherence factor of enteropathogenic *Escherichia coli* with a DNA probe. *J Infect Dis.* 1985;152:560–5.
- Ghilard ACR, Gomes TAT, Elias WP, Trabulsi LR. Virulence factors of *Escherichia coli* strains belonging to serogroups O127 and O142. *Epidemiol Infect.* 2003;131:815–21.
- Gismero-Ordoñez J, Dall'Agnol M, Trabulsi LR, Girón JA. Expression of the bundle-forming pilus by enteropathogenic *Escherichia coli* strains of heterologous serotypes. *J Clin Microbiol.* 2002;40:2291–6.
- Trabulsi LR, Gomes KR, TAT. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg Infect Dis.* 2002;8:508–13.
- Bortoloni M, Trabulsi L, Keller R, Frankel G, Sperandio V. Lack of expression of bundle-forming pili in some clinical isolates of enteropathogenic *Escherichia coli* (EPEC) is due to a conserved large deletion in the *bfp* operon. *FEMS Microbiol Lett.* 1999;179:169–74.
- Okeke IN, Borneman JA, Shin S, Mellies JL, Quinn LE, Kaper JB. Comparative sequence analysis of the plasmid-encoded regulator of enteropathogenic *Escherichia coli* strains. *Infect Immun.* 2001;69:5553–64.
- Baldini MM, Kaper JB, Levine MM, Candy DC, Moon HW. Plasmid-mediated adhesion in enteropathogenic *Escherichia coli*. *J Pediatr Gastroenterol Nutr.* 1983;2:534–9.
- Brinkley C, Burland V, Keller R, Rose DJ, Boutin AT, Klink SA, et al. Nucleotide sequence analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid pMAR7. *Infect Immun.* 2006;74:5408–13.
- Tobe T, Hayashi T, Han C, Schoolnik G, Ohtsubo E, Sasakawa C. Complete DNA sequence and structural analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid. *Infect Immun.* 1999;67:5455–62.
- Gomes TAT, Griffin PM, Ivey C, Trabulsi LR, Ramos SRT. EPEC infections in São Paulo. *Rev Microbiol.* 1996;27:25–33.
- Franken J, Franke S, Schmidt H, Schwarzkopf A, Wieler LH, Baljer G, et al. Nucleotide sequence analysis of enteropathogenic *Escherichia coli* (EPEC) adherence factor probe and development of PCR for rapid detection of EPEC harboring virulence plasmids. *J Clin Microbiol.* 1994;32:2460–3.
- Dulguer MV, Fabbriotti SH, Bando SY, Moreira-Filho CA, Fagundes-Neto U. Atypical enteropathogenic *Escherichia coli* strains: phenotypic and genetic profiling reveals a strong association between enteroaggregative *E. coli* heat-stable enterotoxin and diarrhea. *J Infect Dis.* 2003;188:1685–94.
- Abe CM, Trabulsi LR, Blanco J, Blanco M, Dahbi G, Blanco JE, et al. Virulence features of atypical enteropathogenic *Escherichia coli* identified by the *eae* + EAF-negative *stx*- genetic profile. *Diagn Microbiol Infect Dis.* 2009;64:357–65.
- Gomes TAT, Irino K, Girão VBC, Guth BEC, Vaz TMI, Moreira FC, et al. Emerging enteropathogenic *Escherichia coli* strains? *Emerg Infect Dis.* 2004;10:1851–5.
- Scaletsky ICA, Fabbriotti SH, Aranda KR, Morais MB, Fagundes-Neto U. Comparison of DNA hybridization and PCR assays for detection of putative pathogenic enteroadherent *Escherichia coli*. *J Clin Microbiol.* 2002;40:1254–8.
- Scaletsky ICA, Fabbriotti SH, Silva SO, Morais MB, Fagundes-Neto U. HEp-2-adherent *Escherichia coli* strains associated with acute infantile diarrhea, São Paulo, Brazil. *Emerg Infect Dis.* 2002;8:855–8.
- Scaletsky ICA, Aranda KR, Souza TB, Silva NP, Morais MB. Evidence of pathogenic subgroups among atypical enteropathogenic *Escherichia coli* strains. *J Clin Microbiol.* 2009;47:3756–9.
- Scaletsky IC, Aranda KR, Souza TB, Silva NP. Adherence factors in atypical enteropathogenic *Escherichia coli* strains expressing the localized adherence-like pattern in HEp-2 cells. *J Clin Microbiol.* 2010;48:302–6.

36. Lacher DW, Steinsland H, Blank TE, Donnenberg MS, Whittam TS. Molecular evolution of typical enteropathogenic *Escherichia coli*: clonal analysis by multilocus sequence typing and virulence gene allelic profiling. *J Bacteriol.* 2007;189:342–50.
37. Leverton LQ, Kaper JB. Temporal expression of enteropathogenic *Escherichia coli* virulence genes in an in vitro model of infection. *Infect Immun.* 2005;73:1034–43.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

